


AP Calculus AB-4 / BC-4

The graph of the function  $f$  shown above consists of two line segments. Let  $g$  be the function given by  $g(x) = \int_0^x f(t) dt$ .



- (a) Find  $g(-1)$ ,  $g'(-1)$ , and  $g''(-1)$ .
- (b) For what values of  $x$  in the open interval  $(-2, 2)$  is  $g$  increasing? Explain your reasoning.
- (c) For what values of  $x$  in the open interval  $(-2, 2)$  is the graph of  $g$  concave down? Explain your reasoning.
- (d) On the axes provided, sketch the graph of  $g$  on the closed interval  $[-2, 2]$ .

(a)  $g(-1) = \int_0^{-1} f(t) dt = -\int_{-1}^0 f(t) dt = -\frac{3}{2}$   
 $g'(-1) = f(-1) = 0$   
 $g''(-1) = f'(-1) = 3$

3 { 1:  $g(-1)$   
 1:  $g'(-1)$   
 1:  $g''(-1)$

(b)  $g$  is increasing on  $-1 < x < 1$  because  $g'(x) = f(x) > 0$  on this interval.

2 { 1: interval  
 1: reason


(c) The graph of  $g$  is concave down on  $0 < x < 2$  because  $g''(x) = f'(x) < 0$  on this interval.

2 { 1: interval  
 1: reason

or

because  $g'(x) = f(x)$  is decreasing on this interval.

(d)




2 { 1:  $g(-2) = g(0) = g(2) = 0$   
 1: appropriate increasing/decreasing and concavity behavior  
 < -1 > vertical asymptote

### Question 5

Let  $f$  be a function defined on the closed interval  $[0, 7]$ . The graph of  $f$ , consisting of four line segments, is shown above. Let  $g$  be the

function given by  $g(x) = \int_2^x f(t) dt$ .

- Find  $g(3)$ ,  $g'(3)$ , and  $g''(3)$ .
- Find the average rate of change of  $g$  on the interval  $0 \leq x \leq 3$ .
- For how many values  $c$ , where  $0 < c < 3$ , is  $g'(c)$  equal to the average rate found in part (b)? Explain your reasoning.
- Find the  $x$ -coordinate of each point of inflection of the graph of  $g$  on the interval  $0 < x < 7$ . Justify your answer.



$$\begin{aligned} \text{(a)} \quad g(3) &= \int_2^3 f(t) dt = \frac{1}{2}(4 + 2) = 3 \\ g'(3) &= f(3) = 2 \\ g''(3) &= f'(3) = \frac{0 - 4}{4 - 2} = -2 \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad \frac{g(3) - g(0)}{3} &= \frac{1}{3} \int_0^3 f(t) dt \\ &= \frac{1}{3} \left( \frac{1}{2}(2)(4) + \frac{1}{2}(4 + 2) \right) = \frac{7}{3} \end{aligned}$$

(c) There are two values of  $c$ .  
We need  $\frac{7}{3} = g'(c) = f(c)$

The graph of  $f$  intersects the line  $y = \frac{7}{3}$  at two places between 0 and 3.

(d)  $x = 2$  and  $x = 5$   
because  $g' = f$  changes from increasing to decreasing at  $x = 2$ , and from decreasing to increasing at  $x = 5$ .

$$3 : \begin{cases} 1 : g(3) \\ 1 : g'(3) \\ 1 : g''(3) \end{cases}$$

$$2 : \begin{cases} 1 : g(3) - g(0) = \int_0^3 f(t) dt \\ 1 : \text{answer} \end{cases}$$


$$2 : \begin{cases} 1 : \text{answer of 2} \\ 1 : \text{reason} \end{cases}$$

Note: 1/2 if answer is 1 by MVT

$$2 : \begin{cases} 1 : x = 2 \text{ and } x = 5 \text{ only} \\ 1 : \text{justification} \\ \quad (\text{ignore discussion at } x = 4) \end{cases}$$

### Question 4

The graph of the function  $f$  above consists of three line segments.



(a) Let  $g$  be the function given by  $g(x) = \int_{-4}^x f(t) dt$ .

For each of  $g(-1)$ ,  $g'(-1)$ , and  $g''(-1)$ , find the value or state that it does not exist.

(b) For the function  $g$  defined in part (a), find the  $x$ -coordinate of each point of inflection of the graph of  $g$  on the open interval  $-4 < x < 3$ . Explain your reasoning.

(c) Let  $h$  be the function given by  $h(x) = \int_x^3 f(t) dt$ . Find all values of  $x$  in the closed interval  $-4 \leq x \leq 3$  for which  $h(x) = 0$ .

(d) For the function  $h$  defined in part (c), find all intervals on which  $h$  is decreasing. Explain your reasoning.

(a)  $g(-1) = \int_{-4}^{-1} f(t) dt = -\frac{1}{2}(3)(5) = -\frac{15}{2}$   
 $g'(-1) = f(-1) = -2$   
 $g''(-1)$  does not exist because  $f$  is not differentiable at  $x = -1$ .

3 :  $\begin{cases} 1 : g(-1) \\ 1 : g'(-1) \\ 1 : g''(-1) \end{cases}$

(b)  $x = 1$   
 $g' = f$  changes from increasing to decreasing at  $x = 1$ .

2 :  $\begin{cases} 1 : x = 1 \text{ (only)} \\ 1 : \text{reason} \end{cases}$

(c)  $x = -1, 1, 3$

2 : correct values  
 $\langle -1 \rangle$  each missing or extra value

(d)  $h$  is decreasing on  $[0, 2]$   
 $h' = -f < 0$  when  $f > 0$

2 :  $\begin{cases} 1 : \text{interval} \\ 1 : \text{reason} \end{cases}$