

Material from the previous units will be covered on this test.

---

A graphing calculator is required for some problems or parts of problems.

---

1. A particle moves along the  $y$ -axis so that its velocity  $v$  at time  $t \geq 0$  is given by  $v(t) = 1 - \tan^{-1}(e^t)$ . At time  $t = 0$ , the particle is at  $y = -1$ . (Note:  $\tan^{-1} x = \arctan x$ )
- (a) Find the acceleration of the particle at time  $t = 2$ .
  - (b) Is the speed of the particle increasing or decreasing at time  $t = 2$ ? Give a reason for your answer.
  - (c) Find the time  $t \geq 0$  at which the particle reaches its highest point. Justify your answer.
  - (d) Find the position of the particle at time  $t = 2$ . Is the particle moving toward the origin or away from the origin at time  $t = 2$ ? Justify your answer.
- 

WRITE ALL WORK IN THE EXAM BOOKLET.

GO ON TO THE NEXT PAGE.

Distance $x$ (cm)	0	1	5	6	8
Temperature $T(x)$ ( $^{\circ}\text{C}$ )	100	93	70	62	55

2. A metal wire of length 8 centimeters (cm) is heated at one end. The table above gives selected values of the temperature  $T(x)$ , in degrees Celsius ( $^{\circ}\text{C}$ ), of the wire  $x$  cm from the heated end. The function  $T$  is decreasing and twice differentiable.
- (a) Estimate  $T'(7)$ . Show the work that leads to your answer. Indicate units of measure.
- (b) Write an integral expression in terms of  $T(x)$  for the average temperature of the wire. Estimate the average temperature of the wire using a trapezoidal sum with the four subintervals indicated by the data in the table. Indicate units of measure.
- (c) Find  $\int_0^8 T'(x) dx$ , and indicate units of measure. Explain the meaning of  $\int_0^8 T'(x) dx$  in terms of the temperature of the wire.
- (d) Are the data in the table consistent with the assertion that  $T''(x) > 0$  for every  $x$  in the interval  $0 < x < 8$ ? Explain your answer.
- 

**WRITE ALL WORK IN THE EXAM BOOKLET.**

**GO ON TO THE NEXT PAGE.**

3. Consider the curve given by  $x^2 + 4y^2 = 7 + 3xy$ .

(a) Show that  $\frac{dy}{dx} = \frac{3y - 2x}{8y - 3x}$ .

(b) Show that there is a point  $P$  with  $x$ -coordinate 3 at which the line tangent to the curve at  $P$  is horizontal. Find the  $y$ -coordinate of  $P$ .

(c) Find the value of  $\frac{d^2y}{dx^2}$  at the point  $P$  found in part (b). Does the curve have a local maximum, a local minimum, or neither at the point  $P$ ? Justify your answer.


---

**WRITE ALL WORK IN THE EXAM BOOKLET.**

**END OF PART A OF SECTION II**

**CALCULUS AB**  
**SECTION II, Part B**  
Time – 45 minutes  
Number of problems – 3


No calculator is allowed for these problems.



4. A coffeepot has the shape of a cylinder with radius 5 inches, as shown in the figure above. Let  $h$  be the depth of the coffee in the pot, measured in inches, where  $h$  is a function of  $t$ , measured in seconds. The volume  $V$  of coffee in the pot is changing at the rate of  $-5\pi\sqrt{h}$  cubic inches per second. (The volume  $V$  of a cylinder with radius  $r$  and height  $h$  is  $V = \pi r^2 h$ .)
- (a) Show that  $\frac{dh}{dt} = -\frac{\sqrt{h}}{5}$ .
- (b) Given that  $h = 17$  at time  $t = 0$ , solve the differential equation  $\frac{dh}{dt} = -\frac{\sqrt{h}}{5}$  for  $h$  as a function  $t$ .
- (c) At what time  $t$  is the coffeepot empty?

**WRITE ALL WORK IN THE EXAM BOOKLET.**

**GO ON TO THE NEXT PAGE.**



5. The graph of the function  $f$  shown above consists of a semicircle and three line segments. Let  $g$  be the function given by  $g(x) = \int_{-3}^x f(t) dt$ .
- Find  $g(0)$  and  $g'(0)$ .
  - Find the values of  $x$  in the open interval  $(-5, 4)$  at which  $g$  attains a relative maximum. Justify your answer.
  - Find the absolute minimum value of  $g$  on the closed interval  $[-5, 4]$ . Justify your answer.
  - Find the values of  $x$  in the open interval  $(-5, 4)$  at which the graph of  $g$  has a point of inflection.
- 

**WRITE ALL WORK IN THE EXAM BOOKLET.**

**GO ON TO THE NEXT PAGE.**

6. Let  $f$  be a function defined by  $f(x) = k\sqrt{x} - \ln x$  for  $x > 0$ , where  $k$  is a positive constant.
- (a) Find  $f'(x)$  and  $f''(x)$ .
  - (b) For what value of the constant  $k$  does  $f$  have a critical point at  $x = 1$ ? For this value of  $k$ , determine whether  $f$  has a relative minimum, relative maximum, or neither at  $x = 1$ . Justify your answer.
  - (c) For a certain value of the constant  $k$ , the graph of  $f$  has a point of inflection on the  $x$ -axis. Find this value of  $k$ .
- 

**WRITE ALL WORK IN THE EXAM BOOKLET.**

**END OF EXAM**

$x$	$f(x)$	$f'(x)$	$g(x)$	$g'(x)$
1	6	4	2	5
2	9	2	3	1
3	10	-4	4	2
4	-1	3	6	7

3. The functions  $f$  and  $g$  are differentiable for all real numbers, and  $g$  is strictly increasing. The table above gives values of the functions and their first derivatives at selected values of  $x$ . The function  $h$  is given by  $h(x) = f(g(x)) - 6$ .
- (a) Explain why there must be a value  $r$  for  $1 < r < 3$  such that  $h(r) = -5$ .
- (b) Explain why there must be a value  $c$  for  $1 < c < 3$  such that  $h'(c) = -5$ .
- (c) Let  $w$  be the function given by  $w(x) = \int_1^{g(x)} f(t) dt$ . Find the value of  $w'(3)$ .
- (d) If  $g^{-1}$  is the inverse function of  $g$ , write an equation for the line tangent to the graph of  $y = g^{-1}(x)$  at  $x = 2$ .
-

$t$ (minutes)	0	2	5	7	11	12
$r'(t)$ (feet per minute)	5.7	4.0	2.0	1.2	0.6	0.5

5. The volume of a spherical hot air balloon expands as the air inside the balloon is heated. The radius of the balloon, in feet, is modeled by a twice-differentiable function  $r$  of time  $t$ , where  $t$  is measured in minutes. For  $0 < t < 12$ , the graph of  $r$  is concave down. The table above gives selected values of the rate of change,  $r'(t)$ , of the radius of the balloon over the time interval  $0 \leq t \leq 12$ . The radius of the balloon is 30 feet when  $t = 5$ .

(Note: The volume of a sphere of radius  $r$  is given by  $V = \frac{4}{3}\pi r^3$ .)

- (a) Estimate the radius of the balloon when  $t = 5.4$  using the tangent line approximation at  $t = 5$ . Is your estimate greater than or less than the true value? Give a reason for your answer.
- (b) Find the rate of change of the volume of the balloon with respect to time when  $t = 5$ . Indicate units of measure.
- (c) Use a right Riemann sum with the five subintervals indicated by the data in the table to approximate  $\int_0^{12} r'(t) dt$ . Using correct units, explain the meaning of  $\int_0^{12} r'(t) dt$  in terms of the radius of the balloon.
- (d) Is your approximation in part (c) greater than or less than  $\int_0^{12} r'(t) dt$ ? Give a reason for your answer.
-




---

4. A particle moves along the  $x$ -axis with position at time  $t$  given by  $x(t) = e^{-t} \sin t$  for  $0 \leq t \leq 2\pi$ .

(a) Find the time  $t$  at which the particle is farthest to the left. Justify your answer.

(b) Find the value of the constant  $A$  for which  $x(t)$  satisfies the equation  $Ax''(t) + x'(t) + x(t) = 0$  for  $0 < t < 2\pi$ .

---



5. An oil storage tank has the shape shown above, obtained by revolving the curve  $y = \frac{9}{625}x^4$  from  $x = 0$  to  $x = 5$  about the  $y$ -axis, where  $x$  and  $y$  are measured in feet. Oil flows into the tank at the constant rate of 8 cubic feet per minute.
- Find the volume of the tank. Indicate units of measure.
  - To the nearest minute, how long would it take to fill the tank if the tank was empty initially?
  - Let  $h$  be the depth, in feet, of oil in the tank. How fast is the depth of the oil in the tank increasing when  $h = 4$ ? Indicate units of measure.
-